

## PRODUCT INFORMATION

# WSD CY-GUARD SPRAY ON

## Spray On Sheep Blowfly Protection

### Product Description

**WSD CY-GUARD SPRAY ON** is an external application for preventing blowfly strike in sheep and lambs.

### Type of Preparation

**WSD CY-GUARD SPRAY ON** is a ready-to-use formulation containing 60g/L cyromazine and a scourable dye for topical application.

### Indications

#### Sheep and Lambs:

Prevention for up to 11 weeks but NOT treatment of blowfly strike by *Lucilia cuprina* in sheep.

**Milking Sheep:** Should not be used in sheep producing milk for human consumption.

### Dosage

Always refer to the label. Always apply **WSD CY-GUARD SPRAY ON** undiluted from the original container. It is important to use an approved applicator to deliver a band onto the fleece close to the areas to be protected. There are different application rates and patterns for protection against poll, body and/or crutch strike, and rates also depend on wool length. If a second treatment is required during a season, it should be applied less than 11 weeks after the first treatment with **WSD CY-GUARD SPRAY ON** or less than 14 weeks after treatment with **WSD CY-GUARD SC**.

### Advice On Use

- **WSD CY-GUARD SPRAY ON** is convenient to use but may not provide protection for as long as jetting or spraying with **WSD CY-GUARD SC**.
- **WSD CY-GUARD SPRAY ON** can be used to achieve moderate periods of fly-strike prevention of up to 11 weeks, including where it is not possible to correctly apply **WSD CY-GUARD SC** by dipping or jetting.
- Duration of activity and effectiveness of **WSD CY-GUARD SPRAY ON** depend on correct application technique.
- **WSD CY-GUARD SPRAY ON** must not be diluted before use.
- It is IMPORTANT to use **WSD CY-GUARD SPRAY ON BEFORE** the anticipated onset of a major fly wave. This may be at a particular time of year OR following unseasonal rain or if exceptional blow-fly activity is detected.
- **WSD CY-GUARD SPRAY ON** may be used in sheep with less than 6 weeks wool.
- Consider using **WSD CY-GUARD SPRAY ON** for sheep severely affected by grass seeds. **WSD CY-GUARD SPRAY ON** is also suitable for use in sheep with open wounds provided it is not applied directly onto the wounds.
- Do NOT treat wet sheep. Heavy rain soon after the application of this product may also reduce the period of protection, so avoid use if heavy showers are forecast.
- This product must NOT be mixed with any other product.

# CAUTION

READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY



# WSD Cy-Guard



## SPRAY ON SHEEP BLOWFLY PROTECTION

**Active constituent:**  
60 g/L Cyromazine

When applied as directed will protect long  
wool sheep against fly strike for 11 weeks.

**20 L**



## DIRECTIONS FOR USE RESTRAINTS

DO NOT use on sheep which are producing or may in the future produce milk or milk products for human consumption.

## CONTRAINDICATIONS

This product should not be used on sheep with less than 6 weeks wool as a reduced period of protection may result. This product should not be used on open wounds e.g. those resulting from marking or mulesing operations. This product should not be applied during gusty or windy weather as spray drift will reduce the period of protection.

## DOSAGE & ADMINISTRATION

| Months Since Shearing  | Total Volume/ Sheep (mL) | Volume Per Treatment Band (mL) | No. of Bands | Sheep Treated Per 1L | Critical Comments   |
|--|--------------------------|--------------------------------|--------------|----------------------|---|
| <b>For the prevention of body strike</b>   |                          |                                |              |                      |   |
| Up to 6  | 34                       | 17                             | 2            | 29                   | <ol style="list-style-type: none"> <li>Use a WSD recommended Spray-On applicator gun.</li> <li>Ensure the applicator gun is clean.</li> <li>Calibrate applicator gun using an accurate volume measure.</li> <li>Ensure all areas to be protected are treated.</li> <li>When preventing body strike ensure bands overlap along the midline.</li> <li>This product is not ideal for treating existing strikes. Treated larvae may take 3-4 days to die.</li> <li>Heavy rain following the application of this product could diminish the period of protection.</li> <li>WSD CY-GUARD SPRAY is colored for band identification. This scourable dye will disappear shortly after treatment.</li> </ol>    |
| 7  | 38                       | 19                             | 2            | 26                   |   |
| 8  | 44                       | 22                             | 2            | 22                   |   |
| 9  | 50                       | 25                             | 2            | 20                   |   |
| 10   | 56                       | 28                             | 2            | 17                   |   |
| Lambs less than 15 kg  | 17                       | 17                             | 1            | 58                   |   |
| <b>For the prevention of body and crutch strike</b>  |                          |                                |              |                      |   |
| Up to 6  | 51                       | 17                             | 3            | 19                   | SEE DIAGRAM   |
| 7  | 57                       | 19                             | 3            | 17                   |   |
| 8  | 66                       | 22                             | 3            | 15                   |   |
| 9  | 75                       | 25                             | 3            | 13                   |   |
| 10   | 84                       | 28                             | 3            | 11                   |   |
| Lambs less than 15 kg  | 34                       | 17                             | 2            | 29                   |   |
| <b>FOR THE PREVENTION OF POLL STRIKE</b>   |                          |                                |              |                      |   |
| Rams and sheep susceptible to poll strike should be treated. Apply at least 5 mL of WSD CY-GUARD SPRAY to the areas requiring protection, around the poll and horns. |                          |                                |              |                      |   |

Not to be used for any purpose, or in any manner, contrary to this label unless authorised under appropriate legislation.

## GENERAL DIRECTIONS

DO NOT dilute. DO NOT mix with any other product. Apply directly from the container using WSD recommended spray-on applicator gun. Except for small lambs under 15Kg, the dose volume is determined by the number of months shearing. Apply WSD Cy-Guard according to the diagrams and tables in the Directions for Use. WSD Cy-Guard Spray may be applied to all breeds of sheep, providing protection against flystrike. WSD Cy-Guard Spray is applied as a low-volume treatment onto or near areas where protection is required. When applying, the nozzle of the applicator gun should be approximately 25cm above the fleece, to achieve a 15cm wide band at each pass.

## WITHHOLDING PERIOD/S

**Meat:** DO NOT use less than 7 days before slaughter for human consumption.

**Milk:** DO NOT use on sheep which are producing or may in the future produce milk or milk products for human consumption.

**WOOL HARVEST INTERVAL:** DO NOT use less than 2 months before shearing or fibre collection.

## TRADE ADVICE

Export Slaughter Interval (ESI): Do not use less than 28 days before slaughter for export. Use of this product may result in wool residues that may not comply with European Union environmental quality standards. Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

## SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear rubber gloves. Wash hands after use.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## ADDITIONAL USER SAFETY

Sheep Rehhandling Interval: After treatment wait until sheep are dry before rehandling. Safety Data Sheet (SDS): If additional hazard information is required, refer to the Safety Data Sheet. For a copy phone WSDA (08) 9321 2888 or visit [www.wsdagribusiness.com](http://www.wsdagribusiness.com)

## ENVIRONMENTAL STATEMENTS

DO NOT contaminate dams, rivers or streams with the product or used containers.

## DISPOSAL

Triple or preferably pressure rinse containers and dispose of rinsate in a disposal pit as described below. DO NOT dispose of undiluted chemicals on site. Break crush or puncture and bury empty container in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically marked and setup for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burned.

## STORAGE

Store below 30°C (room temperature). Store in original container tightly closed in a dry, cool place. DO NOT store in direct sunlight. DO NOT re-use the container. Keep out of reach of children.

## CONDITIONS OF SALE

All WSD Agribusiness products are of a high quality, being subjected to rigid quality control. The manufacturer, however, is unable to control the product's subsequent handling or use or the circumstances of each handling or use. This product is sold on the express condition that neither the manufacturer nor the distributor is under any liability for any loss or damage that may arise through such handling or use. If these conditions are unacceptable the container should be returned unopened for credit.



**Manufactured by  
WSD Agribusiness Pty. Ltd.**

ABN 58 009 372 525  
Koojan Avenue, South Guildford WA 6055  
Tel: (08) 9321 2888 Fax: (08) 9479 4088  
[contact@wsdagribusiness.com](mailto:contact@wsdagribusiness.com)  
[www.wsdagribusiness.com](http://www.wsdagribusiness.com)

**APMVA Approval No: 61663/0614**

**Batch:**

**Expiry:**

