

## PRODUCT INFORMATION

# WSD COMMAND

## Pour On Lousicide for Sheep

### Product Description

**WSD COMMAND** is a treatment for body lice of sheep.

### Type of Preparation

**WSD COMMAND** is a pour-on formulation for topical use containing 25g/L triflumron- an insect growth regulator (IGR) and a scourable dye.

### Indications

#### Sheep and Lambs:

Control of biting lice (*Bovicola ovis*) in sheep off-shears and lambs up to 2 months of age

**Milking Sheep:** Do NOT use in sheep producing milk for human consumption

### Dosage

Always refer to the label. The correct dose for the weight range of sheep should be applied as a single continuous strip down the middle of the back between the poll and the butt of the tail. It is important to use an approved applicator or one with a wide band nozzle.

It is recommended that treated sheep be held together in the yards for 2 hours after treatments and not be let straight out of the race into the paddock.

### Advice On Use

- Triflumuron is an insect growth regulator, not a knockdown insecticide. It prevents the development of immature lice present in the fleece at application and those that hatch from eggs in the following 20 weeks.
- **WSD COMMAND** is applied undiluted.
- It must not be stored in any container other than the original including overnight. Do not leave **WSD COMMAND** containers uncovered or in direct sunlight.
- It is critical that sheep be cleanly shorn without residual clumps or tufts of wool and **WSD COMMAND** must be applied within 7 days of shearing (preferably within 24 hours) to ensure effectiveness.
- Correct application is critical and it should not be applied as a single spot or to sheep which are wet.
- Light to moderate rainfall soon after application should not reduce efficacy.
- A clean muster and single shearing with immediate correct treatment of every animal is ideal for achieving effective control of lice.
- Lambs at foot up to 2 months of age should be correctly treated with an adequate dose if the ewes are treated.
- If a split shearing is inevitable it is critical that treated and untreated sheep be kept separate.
- The repeated use of any IGR is likely to result in the emergence of resistance in sheep lice. It is therefore recommended to regularly alternate treatment with a dip such as **WSD FLOCKMASTER MKII**. Research suggests that this should be after 2 successive years of using of any IGR product.

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY



# WSD Command



## POUR ON LOUSICIDE FOR SHEEP

**Active Constituent:**  
**25 g/L Triflumuron**

For the control of Body Lice *Bovicola ovis* on  
shorn sheep up to 7-days off-shears and lambs  
at foot. Insect growth regulator.

**20 L**



**DIRECTIONS FOR USE  
RESTRAINTS**

DO NOT use on female sheep which are producing or may in the future produce milk or milk products for human consumption.  
DO NOT use on unshorn lambs more than 2 months old; on adult sheep more than 7 days off-shears, or which cannot be cleanly shorn for any reasons e.g. mycotic dermatitis.

**GENERAL DIRECTIONS**

Shake well before use. WSD Command is a knockdown insecticide, it is an insect growth regulator. It prevents the development of immature lice present in the fleece at application and those that hatch from eggs in the following twenty weeks. WSD Command produces a rapid decline in lice numbers with all adult lice dying by fourteen weeks. Sheep should be treated dry within seven days of shearing. Moderate rainfall even within four days will not reduce the products efficacy. All sheep in the mob should be treated. As with all lousicides, management practices which contribute to good lice control, such as clean musters, good fencing and quarantining of newly purchased stock must be practiced. Treated sheep should not be mixed with lousy sheep. Except for ewes and their lambs, treated sheep should not be mixed with lice free sheep for at least four weeks. After four weeks, adult lice on treated sheep cannot infect clean sheep. Treated sheep can occasionally develop backline dermatitis following prolonged exposure to extreme heat and sunlight without access to shade. DO NOT leave container, hose or applicator in the sun for long periods of time. WSD Command may be used in conjunction with worm control drenches and at the same time as vaccination. A scourable blue marker dye is included in WSD Command for specific identification of treated animals. WSD Command can be detrimental to certain plastics, paints and surface finishes. Avoid contact with such surfaces.

**DOSAGE & ADMINISTRATION:**

Apply in a single strip down the middle of the back between the poll and the butt of the tail using an applicator with a wide band nozzle. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams and lambs). A representative sample of animals should be weighed before treatment either with scales or weighband. Where there is a large variation in size within the group, dose rates should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate to avoid excessive overdosing.

**Ewes & Lambs:** Unshorn lambs should be treated to bodyweight according to the dosage table. WSD Command controls lice in ewe/lamb mobs in the following situation:

- a) When shearing ewes with lambs at foot (up to 2 months of age) – shear and treat the ewes as above; treat the unshorn lambs at the same time.
- b) When lambing off the board – treat the ewes at shearing; treat the unshorn lambs at any time (e.g. at marking) up to 2 months of age.

Sheep	Dose	20L pack treats
Under 10kg	10mL	2000
11-20kg	12mL	1666
21-30kg	15mL	1333
31-55kg	20mL	1000
55-75kg	25mL	800
75-85kg	30mL	666
85-95kg	35mL	571

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD/S**

**Meat:** DO NOT use less than fourteen (14) days before slaughter for human consumption.

**Milk:** DO NOT use on sheep which are producing or may in the future produce milk or milk products for human consumption.

**WOOL HARVEST INTERVAL: DO NOT use less than 2 months before shearing or fibre collection.**

**TRADE ADVICE: Export Slaughter Interval (ESI)**

DO NOT use slaughter for export within 66 days of treatment. Use of this product may result in wool residues that may not comply with European Union environmental quality standards. Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

**SAFETY DIRECTIONS**

Harmful if inhaled or swallowed. Will irritate eyes and skin. Avoid contact with eyes and skin. DO NOT inhale vapour. When using the product wear elbow-length butyl rubber gloves. If product in eyes, wash out immediately with water. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking wash hands, arms and thoroughly with soap and water. After each days use wash gloves and contaminated clothing.

**FIRST AID**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**ADDITIONAL USER SAFETY**

Material Safety Data Sheet (MSDS): Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

Sheep Rehandling Interval: After treatment wait until sheep are dry before rehandling.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT**

DO NOT contaminate dams, ponds, rivers, waterways or drains with pesticide or used containers.

**DISPOSAL**

Triple rinse containers with diesel, and dispose of rinsate in a disposal pit away from desirable plants and their roots and watercourses. DO NOT dispose of un-diluted chemicals on-site. Break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and setup for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

**STORAGE**

Store below 30°C (room temperature) in the closed, original container in a dry, well-ventilated area and protect from light.

**CONDITIONS OF SALE**

All WSD Agribusiness products are of a high quality, being subjected to rigid quality control. The manufacturer, however, is unable to control the product's subsequent handling or use or the circumstances of each handling or use. This product is sold on the express condition that neither the manufacturer nor the distributor is under any liability for any loss or damage that may arise through such handling or use. If these conditions are unacceptable the container should be returned unopened for credit.

Registered, manufactured and distributed in Australia by



WSD Agribusiness Pty. Ltd.  
Koojan Avenue, South Guildford WA 6055  
Tel: (08) 9321 2888 Fax: (08) 9479 4088  
contact@wsdagribusiness.com  
www.wsdagribusiness.com

**HAZARD STATEMENTS**

Causes skin irritation. Causes serious eye irritation. May damage fertility or the unborn child. May cause respiratory irritation. Toxic to aquatic life

**PRECAUTIONARY STATEMENTS**

Obtain special instructions before use. Use only outdoors or in a well-ventilated area. Use personal protective equipment as required. Avoid breathing dust/ fume/gas/mist/vapours/spray. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection.

APVMA No: 56254/0614

Batch:

Expiry:

