

PRODUCT INFORMATION

WSD FENBENDAZOLE 100

Oral Anthelmintic for Cattle and Horses

Product Description

WSD FENBENDAZOLE 100 is a broad spectrum treatment for roundworm and tapeworm parasites of cattle and horses

Type of Preparation

WSD FENBENDAZOLE 100 is a suspension for oral use as a drench containing 100g/L fenbendazole- a benzimidazole (BZ).

Indications

Cattle:

Mature and immature stages of the following worms.

Barber's pole worm (*Haemonchus spp*), hair worm or small hair worm (*Trichostrongylus spp*), small brown stomach worm (*Ostertagia ostertagi* including hypobiotic larvae), small intestinal worm (*Cooperia*), whipworm (*Trichuris*), lungworm (*Dictyocaulus viviparus*), *Capillaria spp*, hookworm (*Bunostomum sp*). Fenbendazole aids in the control of tapeworms (*Moniezia spp*).

Dairy Cattle:

Suitable for cattle producing milk for human consumption (see withholding period information)

Horses:

Mature and immature large strongyles (*Strongylus spp*), small strongyles (*Cyanthostomidae*), ascarids (*Parascaris equorum*), pinworms (*Oxyuris equi*), *Strongyloides westeri*.

Dosage

Always refer to the label. BZ's generally have low toxicity but to avoid residue problems, it is good practice to separate into different weight classes if there is a large variation in age and weight.

Cattle: Dose at the rate of 3.75ml per 50kg body weight.

Horses: Dose at the rate of 1ml per 10kg body weight

Advice On Use

- **WSD FENBENDAZOLE 100** is a reliable broad-spectrum oral drench for cattle and horses of all ages.
- **WSD FENBENDAZOLE 100** is safe for use in pregnant mares and stud males.
- It is effective in treating severe scouring due to *Ostertagia*- type II ostertagiosis in adult cattle.
- It is often best administered to adult cattle with a drenching hook.
- The repeated use of any drench is likely to result in the emergence of resistance among cattle parasites. WSD provides a range of treatments for cattle parasites and consideration should be given to utilizing more than one treatment group in a worm control program.

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



WSD Fenbendazole 100

**ORAL ANTHELMINTIC FOR CATTLE
AND HORSES**

Active constituent:
100 g/L Fenbendazole

For the control of benzimidazole sensitive
roundworms and lungworms in cattle,
and large and small strongyles ascarids
and pinworms in horses

10 L



WSD FENBENDAZOLE 100 ORAL ANTHELMINTIC FOR CATTLE AND HORSES

Contains Fenbendazole, a member of the benzimidazole family of drenches. It is effective in cattle and horses against benzimidazole sensitive strains of the following internal parasites:

CATTLE

<i>Haemonchus placei</i>	Barber's pole worm
<i>Ostertagia spp</i>	Small brown stomach worm
<i>Trichostrongylus spp</i>	Small hair worm
<i>Oesophagostomum radiatum</i>	Nodule worm
<i>Nematodirus spp</i>	Thin necked intestinal worm
<i>Cooperia spp</i>	Small intestinal worm
<i>Dictyocaulus viviparus</i>	Lungworm
<i>Trichuris spp</i>	Whipworm

HORSES

<i>S. vulgaris, S. edentates, S. equinus</i>	Large strongyles
<i>Trichonema spp, Triodontophorus spp</i>	Small strongyles
<i>Parascaris equorum</i>	Ascarids
<i>Oxyuris spp</i>	

DIRECTIONS FOR USE: Shake well before use.

ADMINISTRATION:

WSD FENBENDAZOLE 100 can be administered orally through standard accurate drenching equipment. Clean equipment thoroughly before and after use.

DOSE RATES:

Dose rates should be based on the heaviest animal by liveweight in each group (bulls, cows, steers and calves); (horses) and these should be weighed with scales or weighband. Do not underdose. A representative sample of animals should be weighed before treatment either with scales or weighband. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

CATTLE: 3.75mL/50kg bodyweight (7.5mg Fenbendazole per kg bodyweight)

HORSES: 1mL/10kg bodyweight (10mg Fenbendazole per kg bodyweight)

For the control of *Strongyloides westeri* in foals: 5mL/10kg bodyweight (50mg-Fenbendazole per kg bodyweight) Internal parasites may develop resistance to drenches (anthelmintics). Ask your local veterinary adviser for recommended parasite management practices for your area to reduce development of resistance.

Bodyweight CATTLE	Dose Rate Bodyweight	10 L Treats	Bodyweight HORSES	Dose Rate Bodyweight	10 L Treats
50kg	3.75mL	2666	100kg	10mL	1000
100kg	7.50mL	1333	150kg	15mL	667
150kg	11.25mL	888	200kg	20mL	500
200kg	15.00mL	666	250kg	25mL	400
250kg	18.75mL	533	300kg	30mL	333
300kg	22.50mL	444	350kg	35mL	286
350kg	26.25mL	380	400kg	40mL	250
400kg	30.00mL	333	450kg	45mL	222
450kg	33.75mL	296	500kg	50mL	200
500kg	37.50mL	266	550kg	55mL	182
550kg	41.25mL	242	600kg	60mL	167
600kg	45.00mL	222	650kg	65mL	154
650kg	48.75mL	205	700kg	70mL	143
700kg	52.5mL	190	750kg	75mL	133
750kg	56.25mL	177			
Over 750kg	3.75mL per 50kg		Over 750kg	1 mL per 10kg	

It is advisable that a drench resistance test be conducted before any drench is used. If drench failure with this preparation is suspected, seek veterinary confirmation and notify the manufacturer.

WITHHOLDING PERIOD:

MEAT (Cattle): DO NOT USE less than fourteen (14) days before slaughter for human consumption.

MEAT (Horses): DO NOT USE less than twenty eight (28) days before slaughter for human consumption.

MILK: NIL

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI):

DO NOT USE less than fourteen (14) days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on www.wsdagribusiness.com or the APVMA website www.apvma.gov.au/residues/ESI.shtml.

SAFETY DIRECTIONS:

Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre, Australia 13 11 26.

MATERIAL SAFETY DATA SHEET (MSDS): Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT:

Dangerous to fish. Do not contaminate streams, rivers and waterways with this product or used containers.

STORAGE:

Store below 30°C (room temperature) in the closed, original container in a dry, well-ventilated area, and protect from light.

DISPOSAL:

Do not dispose of undiluted chemicals on-site. Break, crush, puncture or bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

CONDITIONS OF SALE:

All WSD Agribusiness products are of a high quality, being subjected to rigid quality control. The manufacturer, however, is unable to control the product's subsequent handling or use or the circumstances of each handling or use. This product is sold on the express condition that neither the manufacturer nor the distributor is under any liability for any loss or damage that may arise through such handling or use. If these conditions are unacceptable the container should be returned unopened for credit.

Registered, manufactured and distributed in Australia by:



APVMA No: 40427/10L/0210

Batch:

Expiry:

WSD Agribusiness Pty. Ltd.
Koojan Avenue, South Guildford WA 6055
Tel: (08) 9321 2888 Fax: (08) 9479 4088
contact@wsdagribusiness.com
www.wsdagribusiness.com

